

OBSERVATIONAL EVIDENCE FOR YOUNG RADIO GALAXIES ARE TRIGGERED BY ACCRETION DISK INSTABILITY

QINGWEN WU

Korea Astronomy and Space Science Institute, Daejeon 305348, Republic of Korea; Email: qwwu@shao.ac.cn
ApJ Letter, in press

ABSTRACT

The bolometric luminosities and black hole (BH) masses are estimated by various methods for a sample of young radio galaxies with known ages. We find that the ages are positively correlated with the bolometric luminosities in these young radio galaxies, which is consistent with theoretical prediction based on radiation pressure instability of accretion disk in Czerny et al. The ages of young radio galaxies are also found to be consistent with the theoretical durations of outbursts in BH mass and accretion rate (in Eddington unit) plane, where the outbursts are assumed to be triggered by the radiation pressure instabilities. Our results provide the observational evidence for the radiation pressure instability, which causes limit-cycle behavior, as a physical mechanism that may be responsible for these short-lived young radio galaxies.

Subject headings: galaxies: active, quasars—accretion, accretion disks—black hole physics, instabilities

1. INTRODUCTION

Gigahertz-peaked spectrum (GPS: projected linear size $D \lesssim 1$ kpc) and compact steep-spectrum (CSS: projected linear size $D \lesssim 20$ kpc) radio sources constitute a large fraction ($\sim 40\%$) of the powerful radio source population. Their radio spectra are simple and convex with peaks close to 1 GHz and 100 MHz for GPS and CSS sources respectively (see O’Dea 1998 for a review). Two basic models have been proposed to explain the CSS/GPS phenomena: (1) *youth scenario*: the radio sources are compact because they are young, and the jets will expand into large-scale ‘classical’ lobes during the course of their life (e.g., Fanti et al. 1995); (2) *frustration scenario*: the radio emitting plasma is permanently confined to the region within the host galaxy by a cocoon of dense gas and dust for their entire life (e.g., van Breugel 1984). Both kinematic ages estimated from proper motions of hot spots (e.g., Giroletti & Polatidis 2009 and references therein) and radiative ages deduced by synchrotron theory based on the radio spectra of lobes (e.g., Murgia 1999 and reference therein) provide values from $\sim 10^2$ to $\sim 10^5$ years, which strongly support the *youth scenario*.

Duty cycle is one of key problems in understanding the evolution of active galactic nuclei (AGN, e.g., Wang et al. 2009, and references therein). The extended radio emission in galaxies and quasars provides us with an opportunity to probe their history through the ages derived from the structural and spectral information of their radio lobes. Both the young compact symmetric object (CSO) with a much older extended relic emission (e.g., Owsianik et al. 1998) and the ‘double-double’ radio source which consists of a new lobe-pair near the core and a more distant faded lobe-pair (e.g., Lara et al. 1999; Schoenmakers et al. 2000) can be explained in terms of recurrent activity in their nuclei. These possibly re-born radio sources may be intermittent on timescales of $\sim 10^4 - 10^5$ years (e.g., Reynolds & Begelman 1997).

A standard, geometrically thin accretion disk (SSD, Shakura & Sunyaev 1973) is known to be unstable in the innermost, radiation pressure dominated regions when accretion rate is high (e.g., Lightman & Eardley 1974). The thermally unstable inner region will possibly lead to the so-called limit-cycle behavior, which has been confirmed by numerical simulations of the time-dependent disk (e.g., Honma et al. 1991; Li et

al. 2007). On the observational side, a perfect candidate of black hole (BH) X-ray binary GRS 1915+105 has been known to show the theoretically predicted limit-cyclic luminosity variations (e.g., Belloni et al. 1997; Nayakshin et al. 2000; Janiuk et al. 2002; Merloni & Nayakshin 2006). Czerny et al. (2009) found that the outburst durations are generally from $\sim 10^2$ to $\sim 10^5$ years for BH mass $M_{\text{BH}} = 10^7 - 10^{10} M_{\odot}$ and viscosity parameter $\alpha = 0.02 - 0.2$, which are roughly consistent with the observed ages of young radio galaxies, where the viscosity law with a geometrical mean between the gas and the total pressure for scaling of the stress, i.e., $T_{r\phi} \propto (P_{\text{gas}} P_{\text{tot}})^{1/2}$, is adopted when modeling the time evolution of SSD under the radiation pressure instability. An important empirical correlation between the outburst duration, T_{burst} , bolometric luminosity, L_{bol} , and viscosity parameter, α , is given by Czerny et al. (2009),

$$\log \left(\frac{T_{\text{burst}}}{\text{yr}} \right) \simeq 1.25 \log \left(\frac{L_{\text{bol}}}{\text{erg/s}} \right) + 0.38 \log \left(\frac{\alpha}{0.02} \right) - 53.6. \quad (1)$$

Wu (2009) found that most of young radio galaxies have relatively high accretion rates with average Eddington ratio $< \log L_{\text{bol}}/L_{\text{Edd}} > = -0.56$ for a sample of 51 sources (32 CSS + 19 GPS), where $L_{\text{Edd}} = 1.38 \times 10^{38} M_{\text{BH}}/M_{\odot}$ is Eddington luminosity. The accretion disk in these high Eddington ratio CSS/GPS sources may have suffered the radiation pressure instability, which provide a possibility to explain their short-lived activities. In this *Letter*, we estimate the bolometric luminosities and BH masses for a sample of young radio galaxies with known ages. We then test whether the ages of young radio galaxies are consistent with the durations of outbursts or not, where the outbursts are assumed to be triggered by the radiation pressure instabilities of accretion disk (e.g., Czerny et al. 2009). Throughout this paper, we assume the following cosmology: $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_0 = 0.3$ and $\Omega_{\Lambda} = 0.7$.

2. THE SAMPLE

We compile a sample of 37 young radio sources from the data available in the literature. The kinematic ages are mainly selected from Giroletti & Polatidis (2009, and references therein), while the ages estimated from synchrotron cooling time are mainly taken from Murgia et al. (1999). We give the lower limits of kinematic age for four sources (0134+329, 0538+498,

1458+718, 2249+185) assuming the upper limit for the separation of hot spots to be the light speed c in this work, since their radiative ages may be much shorter than the actual source ages in these jets or hot spots dominated sources (see Murgia et al. 1999 for more details). The kinematic age of CSS source 1328+307 (3C 286) is estimated in the present work from its projected liner size and jet propagation speed $\simeq 0.1c$ (An et al. 2004). Several other individual sources with known ages are also selected from the literature. In total, our sample consists of 14 GPS and 23 CSS sources (see Table 1 for more details).

3. BOLOMETRIC LUMINOSITIES AND BH MASSES

The bolometric luminosity of AGN is sometimes approximated from the optical luminosity, since the integration of spectral energy distribution (SED) is usually hampered by the lack of multi-wavelength observational data that spans many decades in wavelength. It is found that the bolometric luminosity is roughly 9 times the optical luminosity, $L_{\text{bol}} = 9L_{5100}$, where $L_{5100} = \lambda L_{\lambda}(5100)$ (Kaspi et al. 2000). It should be noted that the monochromatic optical luminosity cannot be used to derive the bolometric luminosity if the optical luminosity is contaminated by the jet emission in radio loud AGN (e.g., BL Lacs) or obscured by the putative torus. The broad line emission is believed to be illuminated by the ionizing luminosity from the accretion disk, and, therefore, it can be used as a good indicator of thermal optical emission. We estimate the bolometric luminosities for 9 sources from their broad line luminosities (e.g., $H\beta$, C IV) using the empirical correlations between the luminosities of different emission lines and bolometric luminosities derived from a sample of radio-quiet AGN in Wu (2009). The bolometric luminosities of 2 sources fitted from their SEDs are selected directly (2342+821, 1458+718, Woo & Urry 2002). The bolometric luminosity of 1 source have been corrected for beaming effect is also included (1413+135, Xie et al. 2006). The bolometric luminosities of 4 optically Type 1 CSS sources are estimated from their optical luminosities with $L_{\text{bol}} = 9L_{5100}$, since that, normally, the steep-spectrum sources do not suffer strong contamination from the Doppler-boosted emission of the jet due to the possible large inclination angles.

In the standard model, narrow emission lines (e.g., [O III]) are also believed to be excited through photoionization by the UV continuum produced by the accretion disk (e.g., in radio-quiet AGN). However, it is more complicated in young radio galaxies, since shocks caused by the interaction between the jet and interstellar medium (ISM) may also be important for the morphology, kinematics, and the excitation of narrow emission lines (e.g., Labiano et al. 2005; Holt et al. 2006). Moy & Rocca-Volmerange (2002) proposed that the shock ionization may be only important for sources with intermediate radio sizes ($2 \text{ kpc} < D < 150 \text{ kpc}$, e.g., CSS sources), while AGN photoionization is dominant in the most compact sources ($D < 2 \text{ kpc}$, e.g., GPS sources) and the extended sources with $D > 150 \text{ kpc}$. Moreover, the jet sizes in all GPS sources of our sample are less than 0.2 kpc (see Table 1), which are much less than the kpc-scale of narrow line regions. This also support that the shock contribution to narrow lines from the jet-cloud interaction may be not important and, therefore, narrow line emission may still be a good indicator of central thermal emission in these GPS sources. To derive the bolometric luminosities for more sources, we investigate the relation between [O III] luminosities and bolometric luminosities for a sample of radio-quiet Type 1 AGN, which include 20 Seyferts and 23 low red-shift

QSOs. The Seyferts are selected from Kaspi et al. (2000), where the [O III] luminosities are chosen from Kraemer et al. (2004), and the QSOs are selected from Alonso-Herrero et al. (1997). To be consistent with other methods, we calculate the bolometric luminosities from $L_{\text{bol}} = 9L_{5100}$ for these radio-quiet AGN. The relation between the bolometric luminosities and [O III] luminosities is shown in Figure 1, and its linear regression is,

$$\log L_{\text{bol}} = 0.95 \pm 0.08 \log L_{[\text{O III}]} + 5.39 \pm 3.17. \quad (2)$$

We find that bolometric correction factors, $L_{\text{bol}}/L_{[\text{O III}]}$, are about 2000 and 3400 for 20 Seyferts and 23 QSOs respectively. The bolometric luminosities for 12 GPS and 9 CSS sources were calculated with Eq. (2) from their [O III] luminosities. One should keep in mind that the bolometric luminosities of 9 CSS sources may be over-estimated due to the possible shock contribution to [O III] luminosities (see discussions in Sect. 5).




FIG. 1.— The correlation between bolometric luminosities and [O III] luminosities for a sample of radio-quiet Type 1 AGN. The circles and squares are correspond to 20 Seyferts and 23 QSOs respectively. The solid line is the best fitting.

Wu (2009) have estimated BH masses for a sample of 65 young radio galaxies from different methods. We estimate BH masses of other 14 young radio galaxies, which are not included in Wu (2009), from the relation between host galaxy absolute magnitude M_R in R-band and BH mass proposed by McLure et al. (2004),

$$\log_{10}(M_{\text{BH}}/M_{\odot}) = -0.5M_R - 2.74. \quad (3)$$

4. RESULTS

We estimate the bolometric luminosities for 37 young radio sources by various methods. The relation between ages and bolometric luminosities for young radio galaxies is presented in Figure 2, where the bolometric luminosities of CSS sources calculated from [O III] luminosities are marked with extra open squares. We find that the ages are positively correlated with the bolometric luminosities in these young radio galaxies, which is consistent with the prediction of radiation pressure instability in accretion disk (Czerny et al. 2009). We plot the dashed lines corresponding to Eq. (1) with viscosity parameters $\alpha = 10^{-4}$ and 1 in Figure 2. We find that most sources are locate to the left of the line with $\alpha = 1$ ($L_{\text{bol}} - T_{\text{burst}}$), where the $\alpha = 1$ line gives an upper limit of the theoretical outburst duration. The relation between bolometric luminosity and half of the outburst duration with $\alpha = 0.1$ is also presented in Figure 2 ($L_{\text{bol}} - T_{\text{burst}}/2$, solid line), which is converted from the Eq. (1). We find that the real ages of sources in our sample are roughly consistent with the theoretical prediction of $T_{\text{burst}}/2$ with $\alpha \simeq 0.1$, since

that sources should have the real ages between 0 and T_{burst} (e.g., $0 < \text{age} < T_{\text{burst}}$), and, on the average, the real ages should correspond to $T_{\text{burst}}/2$ with some dispersion around this value.




FIG. 2.— The correlation between ages and bolometric luminosities for young radio galaxies. The dashed lines are theoretical predictions from the disk instability model for $L_{\text{bol}} - T_{\text{burst}}$ with $\alpha = 10^{-4}$ (upper) and 1 (lower) respectively, while the solid line is for $L_{\text{bol}} - T_{\text{burst}}/2$ with $\alpha = 0.1$. The bolometric luminosities of CSS sources calculated from [O III] luminosities are marked with extra open squares.




FIG. 3.— Comparison of the ages for young radio galaxies with the theoretical durations of outbursts triggered by disk instability in $M_{\text{BH}} - \dot{m}$ plane. The triangles, squares, pentagons, and circles correspond to the age in bins of $[0-5 \times 10^2]$, $[5 \times 10^2-5 \times 10^3]$, $[5 \times 10^3-5 \times 10^4]$, and $[5 \times 10^4-5 \times 10^5]$ yrs respectively. The solid lines are the half of theoretical outburst durations predicted by the disk instability model with $\alpha = 0.1$, which are converted from the case of $\alpha = 0.2$ in Czerny et al. (2009, Fig. 5) using the scaling relation of Eq. (1).

The BH masses are calculated from the luminosities of host galaxy, or collected from the literature. The Eddington ratios, $\dot{m} = L_{\text{bol}}/L_{\text{Edd}}$, were computed from the estimated bolometric luminosities and BH masses. We then compare the ages of young radio sources with the half of outburst durations of in BH

mass-accretion rate (in Eddington unit, $M_{\text{BH}} - \dot{m}$) plane, where the outbursts are assumed to be triggered by the radiation pressure instabilities of accretion disk. To do so, we convert the contour map of $\alpha = 0.2$ in Czerny et al. (2009, Fig. 5) to the case of $T_{\text{burst}}/2$ with $\alpha = 0.1$ using the scaling relation of Eq. (1), since we find that most of the young radio sources can, on the average, be described by $\alpha \simeq 0.1$ in the $L_{\text{bol}} - \text{age}$ relation. The ages of young radio sources are divided into four bins as $[0-5 \times 10^2]$, $[5 \times 10^2-5 \times 10^3]$, $[5 \times 10^3-5 \times 10^4]$, and $[5 \times 10^4-5 \times 10^5]$ yrs according to the theoretical lines of 10^2 , 10^3 , 10^4 and 10^5 yrs of Czerny et al. (2009) respectively. The results are shown in Figure 3. We find that, on the average, the ages of young radio sources are also roughly consistent with the half of outburst durations with $\alpha \simeq 0.1$ in $M_{\text{BH}} - \dot{m}$ plane, even with substantial scatter. In particular, there are 12 sources in the bin of $[5 \times 10^3-5 \times 10^4]$ yrs (red-pentagons), which coincide with the theoretical prediction of 10^4 yrs (red-solid line) very well.

5. SUMMARY AND DISCUSSION

We estimate bolometric luminosities and BH masses for a sample of young radio galaxies with known ages. We find a positive correlation between the ages and bolometric luminosities in these young radio sources (see Fig. 2). This positive correlation is roughly consistent with the theoretical prediction based on the radiation pressure instability of accretion disk in Czerny et al. (2009). We find that the real ages of most sources are shorter than the upper limit of outburst duration with $\alpha = 1$, and, on the average, correspond to the half of outburst durations with $\alpha \simeq 0.1$ (see Fig. 2). We further compare the ages with the half of outburst durations in $M_{\text{BH}} - \dot{m}$ plane, and find that the observational data is also roughly consistent with the theoretical prediction with $\alpha \simeq 0.1$ (see Fig. 3). Therefore, our results provide observational evidence for the radiation pressure instability as a physical mechanism that may also be responsible for these short-lived young radio galaxies, beside the perfect application to the brightest Galactic X-ray binary GRS 1915+105.

The bolometric luminosities for young radio galaxies are estimated from their SED fittings, optical luminosities, broad line luminosities, or narrow line luminosities. In these methods, the most controversial one is that derived from the [O III] luminosity, due to the possible shock contribution caused by the jet-ISM interaction. However, the shock contribution may be not important in the GPS sources as suggested by the ionization diagnostics of different narrow emission lines (e.g., Moy & Rocca-Volmerange 2002). Wu (2009) also found that accretion activities may still play an important role in shaping the kinematics of [O III] narrow line in young radio galaxies. The bolometric luminosity calculated from [O III] luminosity is $\log L_{\text{bol}}([\text{O III}]) = 45.12$ for the GPS source 1404+286, which is consistent with that calculated from the broad line luminosity $\log L_{\text{bol}}(\text{H}\beta) = 45.19$ very well, where the [O III] luminosity is selected from Marziani et al. (2003). This also support that the shock contribution to [O III] luminosity may not be important, and the [O III] luminosity can still be regarded as a good indicator of central thermal emission in these small radio-sized GPS sources. Labiano et al. (2005) found that the contribution to narrow line luminosity from the shocked gas is between 30 and 70% for CSS source 3C 303.1, and this ratio is even lower in other two CSS sources (3C 67 and 3C 277.1), based on the ionization diagnostics of different narrow emission lines. Therefore, the bolometric luminosities of 9 CSS sources calculated from the [O III] luminosities may not be over-estimated

too much, which will not change our main conclusion.

The limit-cycle behavior triggered by the radiation pressure instability of accretion disk provide a good explanation to the luminosity variations of X-ray binary GRS 1915+105 (e.g., Nayakshin et al. 2000; Janiuk et al. 2002; Merloni & Nayakshin 2006). Czerny et al. (2009) proposed that the young radio galaxies may have the similar origin, and the unstable disk caused by radiation pressure may be the physical reason for their short-lived radio activities. We note that the unique limit-cycle variability only appears in the brightest X-ray binary GRS 1915+105 when it radiates at high luminosity (e.g., Belloni et al. 1997). Wu (2009) found that most of the young radio galaxies have relatively high Eddington ratios, and the accretion disk may also have suffered the radiation pressure instabilities in these young radio sources. We collect a sample of 37 young radio galaxies, and explore whether these young radio sources are consistent with the predictions of disk instability model in Czerny et al. (2009) or not. We test this in both the L_{bol} -age relation and the ages in $M_{\text{BH}} - \dot{m}$ plane. We find that the real ages of most sources are, indeed, less than the upper limit of the theoretical outburst durations with $\alpha = 1$ (see Fig. 2). The viscosity parameter, α , should be larger than 10^{-4} , since that the real ages of most sources are longer than its predicted outburst durations (see Fig. 2). We find that the real ages of these young radio sources are roughly consistent with the model predictions with $\alpha \simeq 0.1$, if we assume the real ages, on the average, should correspond to the half of outburst durations (see Fig. 2). The real ages of the young radio sources are also found to be consistent with the half of outburst durations with $\alpha \simeq 0.1$ in $M_{\text{BH}} - \dot{m}$

plane (see Fig. 3). It should be noted that there are still uncertainties in estimating BH masses, bolometric luminosities and ages in these young radio galaxies. Moreover, the simplified disk model was used to derive the Eq. (1), where corona and jet are neglected (see Czerny et al. 2009 for more details). With all these uncertainties in mind, it seems that the observational data of young radio galaxies are in good agreement with the theoretical predictions of disk instability model. We note that the disk instability model also predicts a complex behavior of the jet activities depending on the accretion rate and BH mass: (1) continuous jet at low accretion rate where the accretion disk is stable (e.g., $\dot{m} \lesssim 10^{-2}$, low luminosity AGN); (2) intermittent compact jet with short outburst duration ($\lesssim 10^4$ yr) at high accretion rate, where the disk is unstable and the short-lived outburst of central activity cannot escape from host galaxy (e.g., $\dot{m} \gtrsim 10^{-2}$, GPS and CSO sources); (3) intermittent jet with long outburst duration ($\gtrsim 10^4$ yr) at high accretion rate (e.g., $\dot{m} \gtrsim 10^{-2}$, CSS sources), where the disk is unstable and the jet is able to escape from the host galaxy and possibly grow to large-scale radio source (e.g., FR I/FR II, see Czerny et al. 2009 for more details).

We thank the anonymous referee for constructive comments that helped to improve the paper. We also thank Czerny, B., Cao, X. W., Zheng, S.-M., Lu, J.-F., Pariev, V. and Das, S. for their careful reading of the manuscript, valuable comments and suggestions. This work is partly supported by the NSFC (grants number 10703009).

REFERENCES

- Akujor, C. E., Spencer, R. E., Zhang, F. J., Fanti, C., Ludke, E., & Garrington, S. T. 1993, *A&A*, 274, 752
- Alonso-Herrero, A., Ward, M. J., Kotilainen, J. K. 1997, 288, 977
- An, T., Hong, X. -Y., Wang, W. -H. 2004, *ChJ&A*, 4, 28
- Axon, D. J., Capetti, A., Fanti, R., Morganti, R., Robinson, A., & Spencer, R. 2000, *AJ*, 120, 2284
- Belloni, T., Mendez, M., King, A. R., van der Klis, M., & van Paradijs, J. 1997, *ApJ*, 488, 109
- Czerny, B., Siemiginowska, A., Janiuk, A., Nikiel-Wroczyński, B., & Stawarz, L. 2009, *ApJ*, 698, 840
- Fanti, C., Fanti, R., Dallacasa, D., McDonald, A., Schilizzi, R. T., & Spencer, R. E. 2002, *A&A*, 396, 801
- Fanti, C., Fanti, R., Dallacasa, D., Schilizzi, R. T., Spencer, R. E., & Stanghellini, C. 1995, *A&A*, 302, 317
- Giroletti, M., Polatidis, A. 2009, *AN*, 330, 193
- Goncalves, A. C., Serote, R., M. 2004, *A&A*, 413, 97
- Gugliucci, N. E., Taylor, G. B., Peck, A. B., & Giroletti, M. 2005, *ApJ*, 622, 136
- Hirst, P., Jackson, N., Rawlings, S. 2003, *MNRAS*, 346, 1009
- Ho, L. C., Peng, C. Y. 2001, *ApJ*, 555, 650
- Holt, J., Tadhunter, C., Morganti, R., Bellamy, M., Gonzalez Delgado, R. M., Tzioumis, A., & Inskip, K. J. 2006, *MNRAS*, 370, 1633
- Honma, F., Matsumoto, R., & Kato, S. 1991, *PASJ*, 43, 147
- Janiuk, A., Czerny, B., & Siemiginowska, A. 2002, *ApJ*, 576, 908
- Kaspi, S., Smith, P., Netzer, H., Maoz, D., Jannuzi, B., Givon, U. 2000, *ApJ*, 533, 631
- Katz-Stone, D. M., Rudnick, L. 1997, *ApJ*, 479, 258
- Kraemer, S. B., George, I. M., Crenshaw, D. M., Gabel, J. R. 2004, 607, 794
- Kunert-Bajraszewska, M., Marecki, A., Thomasson, P. 2006, *A&A*, 450, 945
- Labiano A., et al. 2005, *A&A*, 436, 493
- Lara, L., et al. 1999, *A&A*, 348, 699
- Lawrence, C., Zucker, J. R., Readhead, A., Unwin, S. C., Pearson, T. J., & Xu, W. 1996, *ApJS*, 107, 541
- Li, S.-L., Xue, L., Lu, J.-F. 2007, *ApJ*, 666, 368
- Lightman, A. P., & Eardley, D. M. 1974, *ApJ*, 187, L1
- Marziani, P., Sulentic, J. W., Zamanov, R., Calvani, M., Dultzin-Hacyan, D., Bachev, R., Zwitter, T. 2003, *ApJS*, 145, 199
- McLure, R. J., et al. 2004, *MNRAS*, 351, 347
- Merloni, A., Nayakshin, S. 2006, *MNRAS*, 372, 728
- Moy, E., Rocca-Volmerange, B. 2002, *A&A*, 383, 46
- Murgia, M., et al. 1999, *A&A*, 345, 769
- Nagai, H., Asada, K., Doi, A., Kamenoi, S., Inoue, M. 2009, *AN*, 330, 161
- Nayakshin, S., Rappaport, S., & Melia, F. 2000, *ApJ*, 535, 798
- O'Dea, C., 1998, *PASP*, 110, 493
- Owsianik, I., Conway, J. E., Polatidis, A. G. 1998, *A&A*, 336, 370
- Peck, A. B., Taylor, G. B., Fassnacht, C. D., Readhead, A., Vermeulen, R. C. 2000, *ApJ*, 534, 104
- Reynolds, C. S., Begelman, M. C. 1997, *ApJ*, 487, 135
- Schoenmakers, A. P., de Bruyn, A. G., Rottgering, H., van der Laan, H., Kaiser, C. R. 2000, *MNRAS*, 315, 371
- Shakura, N. I., & Sunyaev, R. A. 1973, *A&A*, 24, 337
- Snellen, I. A. G., Bremer, M. N., Schilizzi, R. T., Miley, G. K., van Ojik, R. 1996, *MNRAS*, 279, 1294
- Taylor, G. B., Inoue, M., Tabara, H. 1992, *A&A*, 264, 421
- Tyulbashev, S. A., Chernikov, P. A. 2001, *A&A*, 373, 381
- van Breugel, W. J. M., 1984, *IAU Symp.*, 110, 59
- Vink, J., Snellen, I., Mack, K.-H., Schilizzi, R. 2006, *MNRAS*, 367, 928
- Wang, J.-M., et al. 2009, *ApJ*, 697, 141
- Woo, J.-H. & Urry, C. M. 2002, *ApJ*, 579, 530
- Wu, Q. W. 2009, *MNRAS*, in press (arXiv:0905.3663)
- Xie, G. Z., et al. 2006, *ApJ*, 131, 1210
- Xu, C., Livio, M., Baum, S. 1999, *AJ*, 118, 1169
- Zakamska, N. L., et al. 2003, *ApJ*, 126, 2125

Table 1. The data of young radio sources.

Name	Others	z	Size kpc	Age [10^3 yrs]	Method	Refs. ^a	L_{bol} log (ergs/s)	Method	Refs. ^b	M_{bh} log($M_{\odot}$)	Refs. ^c	$L_{\text{bol}}/L_{\text{Edd}}$ log
GPS												
0108+388	...	0.669	0.023	0.40	kin	G09	44.20	L(O III)	L96	7.9	21.6 ¹	-1.82
0710+439	...	0.518	0.088	0.93	kin	P03	45.75	L(O III)	L96	8.4	20.0 ¹	-0.75
1031+567	...	0.450	0.109	1.84	kin	T00	45.02	L(O III)	L96	8.1	20.2 ¹	-1.23
1358+624	4C 62.22	0.431	0.160	2.4	kin	V06	45.10	L(O III)	L96	8.2	19.8 ¹	-1.27
1404+286	OQ 208	0.077	0.007	0.22	kin	L07	45.19	L(H β)	M03	8.7	14.6 ¹	-1.69
1934-638	...	0.183	0.085	1.60	kin	G09	45.60	L(O III)	X99	8.5	17.1 ¹	-1.03
2021+614	...	0.227	0.016	0.37	kin	T00	45.17	L(O III)	L96	8.9	16.8 ¹	-1.91
2352+495	...	0.238	0.117	3.0	kin	P03	44.73	L(O III)	L96	8.4	18.0 ¹	-1.79
J1111+1955	...	0.299	0.070	1.62	kin	G05	45.11	L(O III)	P00	8.5	18.4 ¹	-1.48
0116+31	4C 31.04	0.060	0.071	0.50	kin	G09	44.94	L(O III)	X99	...	...	...
1718-649	NGC 6328	0.014	0.002	0.09	kin	G09	44.10	L(O III)	X99	8.4	11.4 ²	-2.44
0316+413	3C 84	0.018	0.015	0.24	kin	N09	44.60	L(O III)	H01	8.5	W02	-2.04
1413+135	...	0.247	0.031	0.13	kin	G05	44.14	Fitting	X06	8.0	18.9 ¹	-1.95
0035+227	...	0.096	0.022	0.45	kin	G09	44.87	L(O III)	G04	...	...	...
CSS												
0221+276	3C 67	0.310	6.8	51	syn	M99	46.27	L(O III)	H03	8.1	W09	0.03
0404+769	4C 76.03	0.599	0.53	1.1	syn	M99	45.34	L(O III)	L96	8.3	20.3 ¹	-1.11
0740+380	3C 186	1.067	8.2	113	syn	M99	46.80	L(H β)	W09	8.9	W09	-0.30
1019+222	3C 241	1.617	2.8	9.3	syn	M99	46.67	L(H β)	W09	7.8	W09	0.56
1203+645	3C 268.3	0.371	3.9	35.4	syn	M99	46.10	L(O III)	H03	7.8	W09	0.16
1250+568	3C 277.1	0.320	4.4	204	syn	M99	45.90	L(H β)	W09	7.6	W09	-0.14
1416+067	3C 298	1.437	9.1	70	syn	F02	47.64	L5100	W09	10.4	W09	-0.83
1443+77	3C 303.1	0.267	5.0	110	syn	M99	46.35	L(O III)	H03	8.4	W09	-0.19
1447+77	3C 305.1	1.132	9.0	151	syn	M99	47.67	L(O III)	A00	8.9	21.0 ²	0.64
2252+12	3C 455	0.543	13.7	348	syn	M99	46.63	L(O III)	X99	8.5	W09	-0.01
2342+821	...	0.735	0.66	1.3	syn	M99	45.82	Fitting	W02	7.5	W09	0.18
0127+233	3C 43	1.459	9.4	200	syn	F02	46.33	L5100	W09	9.2	W09	-1.01
0134+329	3C 48	0.367	2.1	>10	kin	A1	46.61	L(H β)	W09	9.2	W09	-0.30
0518+165	3C 138	0.759	2.9	50	kin	A93	46.42	L(H β)	W09	8.7	W09	-0.42
0538+498	3C147	0.545	2.4	>3.9	kin	A1	45.95	L(H β)	W09	8.7	W09	-0.90
0758+143	3C 190	1.195	14.1	50	syn	K97	45.85	L(H β)	W09	7.8	W09	-0.13
1328+307	3C 286	0.849	14.2	462	kin	A2	46.55	L5100	W09	8.5	W09	-0.09
1328+254	3C 287	1.055	0.4	55 $\pm$ 45	syn	T01	46.89	L5100	W09	9.6	W09	-0.85
1458+718	3C 309.1	0.905	7.8	>13	kin	A1	46.93	Fitting	W02	9.1	W09	-0.31
2249+185	3C 454	1.757	2.1	>3.4	kin	A1	46.14	L (CIV)	W09	8.6	W09	-0.58
0345+337	3C 93.1	0.243	1.2	35.8	syn	M99	45.60	L(O III)	X99	7.1	W09	0.36
1517+204	3C 318	1.574	7.8	150	syn	T92	45.87	L(O III)	H03	...	...	...
0809+404	4C 40.2	0.551	4.8	100	syn	K06	45.96	L(O III)	Z03	8.8	19.4 ²	-0.95

^aReferences for the radio sizes, the ages of these young radio sources, and the methods for deriving the ages.

^bReferences for the data used to derive the bolometric luminosities.

^cReferences for the BH masses and the apparent magnitude of the host galaxies in R -band used to estimate the BH masses in this work, of which (1) and (2) are selected from Snellen et al. (1996) and *NED* (NASA/IPAC Extragalactic Database; <http://nedwww.ipac.caltech.edu>) respectively.

References. — G09: Giroletti & Polatidis 2009; V06: Vink et al. 2006; G05: Gugliucci et al. 2005; N09: Nagai et al. 2009; M99: Murgia et al. 1999; F02: Fanti et al. 2002; A1: this work assuming the upper limit of light speed; A93: Akujor et al. 1993; K97: Katz-Stone & Rudnick 1997; A2: this work using the jet speed of 0.1c in An et al. 2004; T01: Tyulbashev & Chernikov 2001; T92: Taylor et al. 1992; K06: Kunert-Bajraszewska et al. 2006; L96: Lawrence et al. 1996; M03: Marziani et al. 2003; X99: Xu et al. 1999; P00: Peck et al. 2000; H01: Ho & Peng 2001; X06: Xie et al. 2006; G04: Goncalves & Serote 2004; H03: Hirst et al. 2003; W09: Wu 2009 and references therein; A00: Axon et al. 2000; W02: Woo & Urry 2002; Z03: Zakamska et al. 2003.